
Abundance-Based Trend Results for Columbia Basin Salmon and Steelhead ESUs

Tim Fisher, Fisher Fisheries Ltd.

Tim@FisherFisheries.com

and

Rich Hinrichsen

hinrich@seanet.com

For:

Jim Geiselman - KEWN

Bonneville Power Administration
Division of Environment, Fish, and Wildlife

Policy and Planning Branch
911 NE 11th Street
Portland, OR 97208

November 14, 2006

Table Of Contents

1	Introduction and Executive Summary	1
2	Data	1
3	Methods	2
3.1	Visual Inspection of Charts	2
3.2	Mean Abundance Index	3
3.3	Trends In Abundance Index	3
3.4	Interim Recovery Targets	3
4	Results	5
4.1	Datasets	5
4.2	Test Results and Discussion	5
5	Tables	7
6	Figures	30
7	References	41
8	Appendices	45

1 Introduction and Executive Summary

The Action Agencies (AAs) are exploring alternative methods to monitor and assess the status of Evolutionarily Significant Units (ESUs) of salmon and steelhead listed under the Endangered Species Act (ESA). In addition they want to report on the status of unlisted ESUs in the Columbia River basin. There is concern that data to accurately estimate population growth rate (λ), a central indicator in the National Marine Fisheries Service (NOAA Fisheries) [2000 Federal Columbia River Power System \(FCRPS\) Biological Opinion](#) (BiOp), is not readily available, or is not available in a timely fashion. This paper explores population indicators and tests using data that is generally available within a year of the return of adults to the spawning grounds.

We examined and selected a variety of abundance indices or estimates that appear to be readily monitored and available in a timely manner as candidates for performance indicators. Indicators include dam counts; spawners estimated from redd indices, or redd density depending on the data available for each ESU or geographic spawning aggregation within an ESU. Trends are displayed for listed ESUs, and status tests were conducted. Trend was calculated as the slope of the regression of the abundance index (log transformed) versus time. This measure of trend is used instead of λ , because it does not require age-structure data or the reproductive success of hatchery-born spawners, which is lacking for all spawning aggregations. The test assesses whether the trend is greater at the time of the most recent data than it was during the base-period at the time of the BiOp, which we define as either 1980-2000 (herein referred to as a 20-year base period) or 1990-2000 (10-year base period). Another test compares the geometric mean of annual wild abundance at the time of the most recent data to the 5-year geometric mean at the time of the BiOp (1996-2000). These two tests, along with a simple indicator of whether the population growth trend is positive, appear to be the only tractable assessments that can be performed with readily available data.

Thus far this reconnaissance evaluation of data assembled through 2001 - 2006 indicates that the majority of listed and unlisted ESUs, and their component populations, display positive trends due to a broad-based, sharp increase in returning adults realized in recent years. Also, 15 of the 18 ESUs for which we found post-2000 data exhibit substantial increases in abundance after 2000 as compared to the 10-year base period and 16 of the 18 ESUs show substantial increases as compared to the 20-year base period. The performance tests were positive for most, but not all, spawning aggregations that we examined.

We conclude that these abundance indices and the recommended tests are useful in assessing population status of the majority of the ESUs. These tests are simple, and require no complex models or unsupported assumptions. Furthermore, the data needed to conduct these simple tests are readily available in a timely manner for most listed ESUs and can be regularly monitored. Note, however, that the tests are not predictions of the future, but simply track trends in past abundance.

2 Data

Datasets were selected for each of the 19 Columbia River Basin ESUs, both listed and unlisted ([Table 1](#) and [Figures 1-5](#)). We placed a priority on datasets that had complete information from 1980 through 2004, or, if available, 2005. Three types of trend data were used to conduct the selected tests: dam or trap counts (sometimes adjusted for hatchery fish); redd density (redds / linear mile of stream counted) and spawner abundance from redd counts or run reconstructions (for finer geographic/stock specificity). Dam counts are the only way to directly count all members of an ESU for many ESUs; and even they are not entirely accurate due to passage outside of counting hours, fallback, fish misidentification, inconsistent marking, length-at-age assumptions (used for assigning fish to adults and jacks), and estimates of the hatchery fish proportion of the run. Density of redds averaged over an entire ESU can be compared to dam counts; the relationship for the Snake River spring/summer Chinook ESU, for example, is relatively consistent, and the correlation between the two is statistically significant ([Figure 6](#)). Estimates of spawner

abundance in spawning areas are usually made by adjusting redd counts for hatchery fish on the spawning grounds, adults per redd estimates, length of stream surveyed, and other factors. These estimates are inaccurate at best and can be highly misleading if redd counting methodology changes drastically (i.e. the change from index area spawning surveys to a redd census for spring Chinook in the John Day basin beginning in 1999 makes comparisons to earlier estimates questionable). A few estimates of spawner abundance are made from actual spawner or carcass counts; none of these were used in the current analysis.

We used redd count information to estimate minimum adult escapement for larger spawning aggregations than the datasets we found; i.e. Middle Fork, South Fork, and upper Salmon River. In these cases we used the net adults per redd expansion factors for subbasins or streams from published sources (primarily run reconstructions). These trends were used largely to compare against the interim recovery targets set by NOAA Fisheries. We also used the best available data from different sources for many trends. In many cases the trends we used begin with total escapement estimates (adults and jacks) made from dam counts (i.e. Wind River Chinook escapement = Bonneville Dam count (BON) – The Dalles Dam count (TDA) * (average Wind proportion of escapements between BON and TDA)).¹ Finally, we constructed composite trends from available datasets for many ESUs for which entire ESU estimates of abundance via dam counts are not possible. These trends should also be interpreted with caution, especially when comparing recent (post-1980 or -1990) to historical abundance, since time series for individual stocks that contribute to the composite begin in different years and some series have missing years. In general, stocks were used that had complete time series between 1980 and 2000, with the expectation that data for the previous year will be available for each status evaluation conducted by the AAs. Details on the datasets we used or constructed are presented later in this document.

3 Methods

Methods used to assess population status were taken from the draft report of the West Coast Salmon Biological Review Team ([NOAA Fisheries BRT 2003](#)). The two methods were applied to various indices of abundance, which are described above. None of the methods rely on having age structure data or knowing the relative reproductive success of hatchery-born spawners, which is unavailable for nearly all spawning aggregations of anadromous salmonids in the Columbia Basin. These measures can be applied to indices of wild-born spawner abundance or indices of total spawner abundance. The mean abundance index can be used to as a “safety net” test, a simple comparison of population abundance between the base period and current levels. The trend estimate can be used to assess whether the annual population growth rate is greater in the most recent year of data than in the base period. Note that none of these tests are predictions of future performance of the ESUs but rather measures of pre-and post-BiOp performance.

3.1 Visual Inspection of Charts

We constructed charts of each trend that show the raw numbers (almost always adults or natural origin adults), the 5 year geometric mean, and the interim target, if any was found. Trends can be easily determined by inspecting charts in many cases; this is the only “test” we used that allows comparison of recent abundance (i.e. 1980 or 1990 – most recent year) to historical abundance. These charts should be interpreted with caution since methods and sources change frequently in the historical (pre-1980 or -1990) data, and time series for individual stocks that are incorporated in a composite trend begin in different years. In general, time series for individual stocks in a composite are complete from 1980 through 2000.

Progress toward targets can be interpreted from charts for spawning aggregations that have a target, whether established by NOAA Fisheries or calculated as the sum of NOAA Fisheries targets for a geographic area. Again,

¹ This is a fictional example. See Table 2 for examples of early escapement estimates.

these results should be interpreted with caution since in practically all cases the data are incomplete for the geographic unit for which the target was established. Even for stocks with a relatively restricted geographic range (i.e. the Bear Valley/Elk population of Snake River spring/summer Chinook), it is almost certain that not all redds were counted in any year, since spawning censuses (as opposed to redd counts in known spawning areas) are rarely conducted. Even in cases for which “entire ESU” estimates have been made (i.e. Snake spring/summer and fall Chinook), portions of the ESU have not been enumerated (i.e. escapement in a few minor tributaries below Lower Granite Dam (LGR) for spring/summer Chinook and escapement of mainstem spawners and in the lower reaches of tributaries below LGR for fall Chinook).

3.2 Mean Abundance Index

Average abundance of various abundance indices, are reported as a geometric mean. For the sake of comparison, geometric means were calculated for the years 1996-2000 and the years after 2000. The equation for a geometric mean is $geomean = \sqrt[k]{N_t N_{t+1} \cdots N_{t+k-1}}$, where N_t is the population index at time t, and k represents the number of observations entering the geometric mean calculation. When applied to an index of wild-born spawner abundance, the geometric mean supplies the information needed to conduct population-level safety net test. This test compares the current 5-year geometric mean of annual wild abundance to the 5-year geometric mean between 1996 and 2000.

3.3 Trends In Abundance Index

Trends in abundance indices were calculated to assess whether population abundance tends to be increasing, decreasing or staying the same. For comparison, four trends were calculated for each abundance index: the trend based on 1990-2000 (10-year base period), the trend based on 1990-recent data, and trends based on 1980-2000 (20-year base period) and 1980- recent data. In some cases, data were not available for 2004 or '05. In that case, trends were based on data through '02 or '03.

Trend was calculated as the slope of the regression of the abundance index (log transformed) versus time. One was added to the abundance index before log transforming the data to avoid taking the log of zero, which is undefined. Trend was reported as the exponential function of the estimated slope of the regression line. A trend greater than one indicates population increase, a trend less than one indicates population decrease, and a trend of 1 indicates that, on average, population numbers are not changing. The regression equation used was $\ln(N_t + 1) = \beta_0 + \beta_1 t + \varepsilon_t$, where N_t is the abundance index at time t, β_0 is the intercept regression parameter, β_1 is the slope regression parameter, and ε_t is the random error term of the regression. The regression parameter estimates, $\hat{\beta}_0$ and $\hat{\beta}_1$ are obtained through a least squares fit to the data. The trend estimate is then defined as: trend estimate = $\exp(\hat{\beta}_1)$. We stop short of developing confidence intervals for $\exp(\hat{\beta}_1)$ until it can be verified that the residuals are independent and identically distributed.

This measure of trend is used instead of lambda, because it does not require age-structure data (unknown for the majority of the ESUs) or the reproductive success of hatchery-born spawners, which is unknown for all the ESUs. The test assesses whether the trend is greater at the current time than it was during the base-period, which we define as either 1980 or 1990 - 2000.

3.4 Interim Recovery Targets

Interim recovery targets from [Appendix A](#) are presented in [Table 4](#) and [Figure 7](#) for reference purposes only. NOAA Fisheries established targets for the interior Columbia basin ESUs only; we did not find any targets for the Lower

Columbia and Upper Willamette ESUs. We did not conduct nor do we propose any tests to measure progress toward these targets, since these tests would involve complex models and many more assumptions than the currently proposed tests. Where targets were not specified for an entire ESU, targets for individual stocks were summed to produce a target for the ESU, and are presented in [Table 3](#), along with the data (or lack thereof) we have in hand to show progress against the targets. Whether or not this is appropriate is in question, since the BRT did not document their methodology in the source document.

4 Results

4.1 Datasets

We found 109 datasets or spawning aggregations that could be used to calculate trend statistics for all the NOAA Fisheries ESUs, both listed and unlisted. Datasets that we used were complete from at least 1980 through 2000. Some datasets are incomplete for the post-BiOp period (2001-recent) but are nonetheless presented in [Table 4](#). More datasets existed for the more intensely studied ESUs (i.e. SR spring/summer Chinook) than the less intensely studied ESUs; very few datasets were found for the unlisted ESUs. In general the methodology for the [Beamesderfer et al. 1997](#) “index stocks”, Willamette-Lower Columbia Technical Recovery Team (WLC TRT), and Ecosystem Diagnosis and Treatment Model (EDT) processes, redd counts, and recently developed datasets (see [Appendix E](#)) was well documented; we have practically no descriptions of the methodology used to calculate the BRT datasets. Details on the datasets, sources, and their interpretation can be found in [Table 2](#) and Appendices B through E.

Datasets were obtained from the NOAA Fisheries Biological Recovery Team (BRT 2003; [Appendix B](#)), the NOAA Fisheries Willamette/Lower Columbia Technical Recovery Team ([Appendix C](#)), the Northwest Power Planning Council (NPPC) Fish and Wildlife Program (F&WP) contract to construct run reconstructions for the Ecosystem Diagnosis and Treatment model (EDT; [Appendix D](#)), and redd surveys and escapement estimates from various sources ([Appendix E](#)). The spawning aggregations used for the tests are shown in [Appendix F](#). 97 datasets were found that lent themselves to testing (i.e. the data appeared to be reasonable and the period from 1990-2000 was relatively complete). Some datasets were found that estimated abundance for the same spawning aggregation (i.e. [Beamesderfer et al. 1997](#) “index stock” run reconstructions and the BRT interpretation of such); in these cases we used the original source if the time series were of identical length or the longer time series if they were not.

We performed the tests on two classes of datasets: datasets for spawning aggregations which provide the best representation of the population status of each ESU, and datasets for selected, discrete individual populations that partially comprised the ESU datasets. In most cases the datasets that best represented the ESUs were dam counts or estimates of wild spawners made from dam counts; in some cases we used a composite trend of more than one spawning population when no dam counts were available. Many of the individual population datasets were supplied by the BRT.

4.2 Test Results and Discussion

Charts of the 19 ESU trends we used for the tests are shown in [Figure 7](#). Charts of each of the remaining individual populations we tested are presented in [Appendix G](#). Interim targets are included on the charts for reference where they were supplied by NOAA Fisheries or calculated for larger spawning aggregates ([Table 3](#)). Comparison of the geometric mean with the interim recovery target ([Table 4](#)) demonstrates that none of the seven composite trends for which we found targets meet the target in the base period (1990-2000), and that only SR Fall Chinook meet the target in the post-BiOp period.

Of the individual population trends with interim targets, none meet the individual spawning aggregation target during the base period ([Table 6](#)). We note that many of these trends include hatchery fish and all of the targets exclude them, so it is not possible to know whether or not trends that include hatchery adults meet the targets. Inspection of the post-BiOp geometric means (where available), reveals that only Deschutes River steelhead (mid-Columbia ESU), Bear Valley Creek and Imnaha River spring Chinook (Snake River ESU), and Asotin Creek and Joseph Creek steelhead (Grande Ronde; both Snake River ESU) pass the test during the post- BiOp period. Based on other trends we did not test in this analysis, Wenatchee and Entiat River steelhead (Upper Columbia ESU) also meet the interim target during the post-BiOp period.

Two population-level tests were used in this analysis. The first test uses the BRT trend estimate to assess whether the BRT trend estimate is greater at the current time than the base-period of the trend in 2000. For this analysis, the base-period trend was calculated over 1980-2000 and 1990-2000, and the current trend was calculated over 1980- or 1990-2003, 2004, or 2005. In some cases, data were not available in 2004 or '05; in those cases we used data from 1990-2001 or 2002. The test is passed if the trend at the current time exceeds the trend over the base period. We also tested whether the BRT trend is positive (i.e. greater than 1). A third test is a safety net test, which is a simple comparison of the abundance index at the current time to the abundance index in 2000. Specifically, the test compares the geometric mean abundance index for years after 2000, to the geometric mean abundance index during 1996-2000. If the geometric mean at the current time exceeds that geometric mean over 1996-2000, then the test is passed.

Test results are presented in [Tables 4, 5, 6, and 7](#). Results are grouped by ESU and spawning aggregation. Results for the composite trends of each of the 18 ESUs for which we found post-2000 data are displayed in Figures 8 through 10. The results of these tests are illustrated using the percent change in the geometric mean ([Tables 4 and 6; Figure 8](#)), and the percent change in the BRT trends using both a 10 year base period ([Tables 4 and 6; Figure 9](#)) and a 20 year base period ([Tables 5 and 7; Figure 10](#)). When the percent change is positive, the test is passed; otherwise it is not ([Tables 4, 5, 6, and 7](#)). We also examined whether the BRT trend at the current time was positive ("Trend > 1?" in [Tables 4, 5, 6, and 7](#)).

For the 18 composite trends, one for each ESU, we found that the change in mean abundance was almost always passed. The exceptions were LCR chum and LCR steelhead; however many trends in the LCR steelhead ESU composite had missing data for 2001 and none had data after 2001; and the major population of LCR chum (Grays River) is missing for 2004-05, so these results will likely change with more current data. The change in the BRT trend was positive for all ESUs except LCR steelhead, LCR chum, and SR sockeye. Using a 20-year base period did not significantly change the results ([Table 5](#)). Only Snake River sockeye showed a change in the trend tests, from failing to passing the change in BRT trend test.

Of the 68 individual populations we tested, ([Tables 6 and 7](#)), we found that the change in mean abundance was almost always passed; the lone exception was Grande Ronde upper mainstem steelhead (SR ESU). The change in the BRT trend was positive in all except 10 populations. Using a 20-year base period did significantly change the results for the individual populations we tested ([Table 7](#)). Two additional populations, Umatilla River and White Salmon River spring Chinook, had a negative change in the BRT trend, and 18 additional populations had trends that were declining based on a 20-year base period.

In general, the spawning aggregates are the best representation of ESU abundance for stocks which are not a significant component of ocean and freshwater anadromous fisheries. For stocks which experience significant harvest in these fisheries (i.e. greater than 20% of the adult population), a better representation of total population abundance would be the total contribution to the fisheries plus spawning escapement. These stocks include most fall (ocean-type) Chinook, which have experienced significant harvest in the base period. Unfortunately, a full accounting (i.e., run reconstruction) of most of the ESUs that experience high harvest rates is not available at this time. Also, the majority of the lower river ESUs (i.e., those that spawn below Bonneville Dam) have weak or even declining populations, e.g. LCR chum and steelhead. These ESUs are the least affected by the FCRPS. Given these pattern in trends among ESUs that spawn above and below Bonneville Dam, one might theorize that marine and/or freshwater conditions not influenced by the FCRPS as recently configured are a more important factor in limiting population growth than passage through the FCRPS. However, this pattern in trends needs further testing and validation, namely comparison of trends between stocks within and outside of the Columbia River basin. Unfortunately, similar data for most species of Pacific salmon and steelhead are lacking outside the basin.

5 Tables

Table 1. Data description overview. Bold listing status indicates species is listed as either threatened or endangered under the ESA.

ESU	Species	Listing Status	Dam #s	# Spawners	Redd indices	Pop. estimates	Productivity functions
Columbia River	Chum	T	BON (Upper Gorge pop.)	Grays/Chinook, Washougal, Lower Gorge - redds; upper gorge - BON	Grays and Hamilton/Hardy only	N	R/S-BRT, Feb '03
Deschutes River Summer/Fall-run	Chinook	NW	N	TRT Run Recon. (incl. hatch.)	Y (StreamNet)	PSC Chinook TC Deschutes Falls	N?
Lake Wenatchee	Sockeye	NW	RRH, RIS	RIS – RRH (turnoff to Wenatchee R.) - WDFW	N	N?	N?
Lower Columbia River	Chinook	T	Clack., Sandy, Hood	BRT 16 pops.	9 Trends (StreamNet)	PSC Chinook TC Col. Lewis brights	R/S-BRT (10 pops)
Lower Columbia River	Coho	C	Clack., Sandy	3 trends (Busby et al. 1996)	3 trends (Clack., Sandy)	N?	Clack. (Cramer & Cramer)
Lower Columbia River	Steelhead	T	N	Σ 11 select pops. (BRT)	Sandy	N	R/S – BRT 7 pops.
Middle Columbia River	Steelhead	T	N	Σ 4 select pops. (BRT)	6 Σ over subbasin (w/in John Day)	N	N
Middle Columbia River Spring-run	Chinook	NW	Prosser, 3 Mile	9 pops. - TRT Run Recon. (incl. hatch.)	9 trends (Myers et al.)	PSC Chinook TC Col. Upriver Spring	R/S – TRT 10/02 (John Day, Yakima)
Okanogan River	Sockeye	NW	WEL	Expanded spawning ground counts - BC Fisheries?	Y	N?	N?
Snake River	Sockeye	E	ICH-LGR	Redfish L Weir; Sawtooth Weir (recent)	N	N?	N
Snake River Basin	Steelhead	T	ICH-LGR	Σ 9 pops. Incl. LGR aggregate (BRT)	Imnaha	TAC, A-B run only (Al where is this?)	A. Byrne in press
Snake River Fall-run	Chinook	T	ICH-LGR	BRT nat. origin above LGR	Yes – IDFG/NPT/ODFW	N	R/S-BRT, 02/03
Snake River Spring/Summer-run	Chinook	T	ICH-LGR	Nat. sp & su @ LGR; Σ 14 index pops. (BRT); Tucannon – TRT Run Recon.	Yes – IDFG/NPT/SBT	PSC Chinook TC Col. Upriver Spring	R/S-BRT, 02/03; TRT 10/02 (Up, MF, SF Salmon);
Upper Columbia River	Steelhead	E	WEL	Wenatchee/Entiat + Methow/Okanogan	Methow	N	R/S-BRT, Feb '03
Upper Columbia River Spring-run	Chinook	E	PRD-WEL	3 pops. – Wenatchee, Methow, Entiat	9 trends	PSC Chinook TC Col. Upriver Spring	R/S-BRT (3 pops), 02/03, WEN/ENT index
Upper Columbia River Summer/Fall-run	Chinook	NW	PRD-WEL	8 trends (Myers et al.)	5 trends (Myers et al.)	PSC Chinook TC Col. Upriver Summer & URB	N?
Upper Willamette River	Chinook	T	WILF	BRT – Clack., N Santiam, McKenzie	3-4 Trends (StreamNet)	N	N
Upper Willamette River	Steelhead	T	WILF, Foster, Minto	Σ 9 select pops. (BRT)	4 trends (BRT)	N	N

Table 2. Detailed description of information sources, methods, and assumptions used to estimate the adult or total (adult and jack) spawner return run to selected Columbia and Snake River tributaries.

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Lower Columbia Spring Chinook		
Wind River EDT Stock²		
Total run to the river		
1960-1965	Beamesderfer et al. 1997 in Marmorek et al. 1998 .	Original source or methods unknown.
1966-1969	ODFW and WDFW, 2000 (for passage counts at Bonneville Dam); Petit, 2002 (for 1970-1977 Wind River run to the river estimates).	Average of the proportion: Wind River run divided by (Bonneville Dam count less Zone 6 harvest) for years 1970-1977 (equals 0.053), multiplied by each year's Bonneville Dam count for 1966-1969.
Jack and adult run to the river		
1970-1978	Petit, 2002.	WDFW agency database. 2001 estimate is preliminary.
1979-2004	Columbia River Compact 2005	Table A2. Adult spring chinook Zone 6 tributary returns, 1979-2004. Methods largely unknown but comparable to Petit 2002.
Little White Salmon River EDT Stock		
Jack and adult run to the river		
1970-1978	Petit, 2002.	WDFW agency database. 2001 estimate is preliminary.
1979-2004	Columbia River Compact 2005	Table A2. Adult spring chinook Zone 6 tributary returns, 1979-2004. Methods largely unknown but comparable to Petit 2002.
White Salmon River EDT Stock		
Jack and adult run to the river		
1987-2004	Columbia River Compact 2005	Table A2. Adult spring chinook Zone 6 tributary returns, 1979-2004. Methods largely unknown but comparable to Petit 2002.
Hood River BRT Stock		
Total run to the river		
Pre-1963	None.	No escapement estimates are available; adult monthly salmon catch estimates summed over March through July for each year during 1956-1962 range from 0 to 29 (based on angler-returned harvest cards for steelhead and salmon, adjusted for non-response bias). Reported catches could include species or runs other than spring Chinook salmon.
1963-1971	ODFW and CTWSRO 1990 (for dam passage counts); Unpublished sport harvest statistics, ODFW Fish Division.	Sum of Powerdale Dam passage counts and sport harvest of adult salmon (adjusted sums of monthly catches as noted above).
1972-1989	None.	No escapement estimates are available; sport catches of adult spring Chinook salmon during years with open seasons ranged from 0 to 41 (based on annual sport catch statistics from angler-returned harvest cards for steelhead and salmon).

² See attached documentation of [EDT Task 1. Development of Spring and Summer Chinook Biological Information for EDT Validation](#)

Tributary			
Attribute			
Time period	Information sources	Methods, assumptions, or comments	
Hood River BRT Stock			
Adult return to the river			
1989-1991	Beamesderfer et al. 1997 in Marmorek et al. 1998 ; U.S. versus Oregon Technical Advisory Committee, 1997.	The average ratio of the estimated sport catch (based on punch card returns) below Powerdale Dam to Powerdale Dam counts for 1992-1994 (1.33) is used to expand below Powerdale sport catch estimates in 1989-1991 for an escapement estimate. Powerdale Dam passage was not enumerated during those years.	
Jack and adult return to the river			
1992-1998	ODFW, 2001; Unpublished sport harvest statistics, ODFW Fish Division; Olsen et al. 1994; Olsen and French, 1996; Olsen and French, 1998.	Sum of Powerdale Dam passage counts and sport harvest estimates. For years with no creel surveys (1992-1995,1997), sport catch statistics from angler-returned harvest cards are used to estimate adult harvest. Jack dam counts are expanded to include harvest, assuming adult catch rates are the same as jack catch rates.	
1999-2003	Personal communication, 11/20/2001, E. Olsen, ODFW, The Dalles, updated by pers. comm. 8/2004	Sum of Powerdale Dam passage counts and sport harvest creel estimates. (2001 estimate is preliminary).	
WLC TRT ³ Spring Chinook Stocks			
Cowlitz, Kalama, Lewis, and Sandy Rivers – see attached Willamette/Lower Columbia TRT documentation			
WLC TRT Fall Chinook Stocks			
Fourteen Early, Mid- and Late Fall Run Stocks (Sandy River Early Fall; Big White Salmon, Clackamas, Coweeman, Cowlitz, East Fork Lewis, Elochoman, Grays, Kalama, Mill Creek, Washougal, Wind, Fall; Lewis, Sandy Late Fall) – see attached Willamette/Lower Columbia TRT documentation			
Lower Columbia Chum			
WLC TRT Chum stocks			
Hamilton and Hardy Creeks			
Adult spawners			
1967-2000	Willamette/Lower Columbia TRT	Total adults - see attached Willamette/Lower Columbia TRT documentation ; one stock not used since it overlaps the Hamilton/Hardy stock.	
2001-2003	Personal communication, 8/11/2003, Nancy Uusitalo, USFWS Columbia River Office, Vancouver WA, updated by , S. Lorr, USFWS Vancouver WA, pers. comm. 10/2004	Total adults (expansion from peak index count)	
Grays River			
Adult spawners			
1959-2001	Lower Columbia Washington Run Reconstructions	Total natural adult escapement to mainstem and tributaries	
2002-2003	Grays River Chum HGMP	Peak spawners (2003 missing)	
Chinook River			
Adults			
2002-2003	Grays River Chum HGMP	Approximate return to Sea Resources hatchery trap (reintroduced population)	

³ Willamette River/Lower Columbia Technical Recovery Team URL: http://www.nwfsc.noaa.gov/cbd/trt/viability_salmoniddata.htm

PRELIMINARY ABUNDANCE-BASED TREND RESULTS FOR COLUMBIA BASIN SALMON AND STEELHEAD ESUS

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Lower Columbia Steelhead		
WLC TRT Winter Run Steelhead Stocks		
East Fork Lewis, Hood, Kalama, Washougal, Wind Summer; Coweeman, East Fork Lewis, Hood, Kalama, North Fork Toutle, Sandy, South Fork Toutle, Washougal Winter run total adults - TRT dataset summer + winter run stocks; time series start & stop almost at random; use numbers with caution. Generally complete from 1990 – 2000; see attached Willamette/Lower Columbia TRT documentation		
Lower Columbia Coho		
BRT coho stocks		
Adults		
1957-2003	D. Cramer, Doug Cramer@pgn.com , Portland General Electric (PGE), Portland OR, pers. comm. 6/2003, updated by pers. comm.. 5/2004	Adult counts at North Fork Dam (Clackamas River) and Marmot Dam (Sandy River)
2004-2005	Portland General Electric (PGE) 2006	Incomplete until 11/15; not used. Adult counts at North Fork Dam (Clackamas River) and Marmot Dam (Sandy River)
Upper Willamette Spring Chinook		
WLC TRT Spring Chinook Stocks		
Clackamas River Adults		
1958-2002	D. Cramer, Doug Cramer@pgn.com , Portland General Electric (PGE), Portland OR, pers. comm. 5/2003	North Fork dam total adults
2003-2005	Portland General Electric (PGE) 2006	North Fork dam total adults
McKenzie River Adults		
1970-2001	Leaburg dam counts; see attached Willamette/Lower Columbia TRT documentation	Leaburg dam total adults
2002-2003	Personal communication, 8/5/2003, J. Ziller, ODFW, Springfield OR, updated by J. Firman, ODFW, julie.firman@oregonstate.edu , pers. comm. 7/8/2004	Leaburg dam total adults
2004-2005	ODFW 2006	Leaburg dam total adults; no count was made in 2006
Upper Willamette Steelhead		
Willamette Falls Dam Winter Steelhead Stock		
Willamette River @ Willamette Falls (Sullivan Dam) Winter Adults		
1950-2001	D. Domina, PGE, pers. comm.. 7/31/2003	Willamette Falls (Sullivan Dam fish ladder) total winter adult count
2002-2003	J. Firman, ODFW, julie.firman@oregonstate.edu , Pers. Comm.. 7/8/2004	Willamette Falls (Sullivan Dam fish ladder) total winter adult count

PRELIMINARY ABUNDANCE-BASED TREND RESULTS FOR COLUMBIA BASIN SALMON AND STEELHEAD ESUS

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Willamette Falls Dam Winter Steelhead Stock		
2004-2006	ODFW 2006	Willamette Falls (Sullivan Dam fish ladder) total winter adult count
Deschutes River Summer/Fall Chinook Stock		
ESU Totals		
Total run to the river		
1977-2003	S. Pribyl, steve.pribyl@state.or.us , ODFW , The Dalles, pers. comm. 5/2004	Run size of adult fall Chinook salmon in the Deschutes River from Table 29
Mid- Columbia Spring Chinook		
Klickitat River BRT Stock		
Total run to the river		
1954-1965	ODFW and WDFW, 2000 (for passage counts at Bonneville Dam); Update and revisions to Beamesderfer et al. 1997 in Marmorek et al. 1998 , provided by O. Langness, WDFW, 6/6/2000 (for 1966-1976 run to the river estimates); and Petit, 2002 (for 1977 Klickitat River run to the river estimate).	Average of the proportion: Klickitat River run divided by (Bonneville Dam count less Zone 6 harvest) for years 1966-1977 (equals 0.021), multiplied by each year's Bonneville Dam count for 1954-1965.
1966-1976	Update and revisions to Beamesderfer et al. 1997 in Marmorek et al. 1998 , provided by O. Langness, WDFW, 6/6/2000.	Update and revisions are completion products for Bonneville Power Administration Intergovernmental Agreement 96BP90134.
Jack and adult return to the river		
1977-1978	Petit 2002	WDFW agency database. Two-year-old precocious males reported in Petit (2002) are not included in this summary. 2001 estimate is preliminary.
1979-2004	Columbia River Compact 2005	Table A2. Adult spring chinook Zone 6 tributary returns, 1979-2004. Methods largely unknown but comparable to Petit 2002.
Deschutes River Spring Stock		
Total run to the river		
1977-2003	S. Pribyl, steve.pribyl@state.or.us , ODFW , The Dalles, pers. comm. 5/2004	Wild adults trapped at Pelton and Warm Springs hatcheries from Table 14 + wild adult harvest from Table 16

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
John Day River BRT Stock		
Total run to the river (through 1958)		
1954-1958	<p>ODFW and WDFW, 2000 (for passage counts at Bonneville Dam); Beamesderfer et al. 1997 in Marmorek et al. 1998 (for methods for a run reconstruction for John Day Basin spring Chinook salmon); Run reconstruction for John Day spring Chinook salmon prepared for Northwest Power Planning Council as part of an EDT validation project, in progress. Contract C2002-021 between Pacific Northwest Electric Power and Conservation Planning Council and ODFW.</p>	Average of the proportion: John Day River run divided by (Bonneville Dam count less Zone 6 harvest) for years 1959-1967 (equals 0.026), multiplied by each year's Bonneville Dam count for 1954-1958.
Jack and adult run to the river (adults used from 1959-2001)		
1959-1978	Run reconstruction for John Day spring Chinook salmon prepared for Northwest Power Planning Council as part of an EDT validation project , in progress. Contract C2002-021 between Pacific Northwest Electric Power and Conservation Planning Council and ODFW.	A detailed description is presented separately. The run to the river estimate is based on index redd counts in three key spawning areas expanded by area and fish per redd. The estimate includes in-river harvest and pre-spawning mortality. Surveys from 1999-2002 are more intensive than in the past so the redd counts may be inflated, although this is not noted in the documentation supplied by the BRT.
1979-2004	Columbia River Compact 2005	Table A2. Adult spring chinook Zone 6 tributary returns, 1979-2004. Methods largely unknown but comparable to EDT validation run reconstructions.
Redd density		
1959-2003	Carmichael et al. 2001 , updated for 2001-2003 by pers. comm. W. Wilson, wwilson@centurytel.net , ODFW John Day, 5/2003 & 8/2004	Used index area redd counts expanded by a constant adults/redd factor. Surveys from 1999-2002 are more intensive than in the past so the redd density may be inflated.
Umatilla River BRT Stock		
Jack and adult run to the river (used adults)		
1988-1989	CTUIR and ODFW, 1990a.	Hatchery jacks and adults trapped at Threemile Dam. Natural-origin spring Chinook salmon were extirpated from the basin. Re-introduced fish from hatchery production first began to return as adults in 1988.
1990-2001	Draft Umatilla Subbasin/Willow Creek Subbasin Summary, 8/3/2001, prepared for NWPPC., updated by D. Chess, ODFW, Hermiston, OR, pers. comm. 12/3/2001	Threemile Dam passage counts plus harvest downstream of the dam.
Walla Walla River BRT Stock		
Jack and adult run to the river (used adults)		
2000-2001	CTUIR and ODFW, 1990b; Draft Walla Walla Subbasin Summary, 8/3/2001, prepared for NWPPC; personal communication, P. Kissner, CTUIR, 4/10/2002.	Natural-origin spring Chinook salmon were extirpated from the basin. Adults and jacks trapped at Ringgold Springs Hatchery were outplanted to the Walla Walla Basin during the 2000 and 2001 return years. These outplants are assumed to be accounted for in the Ringgold hatchery returns (see below) and are not added into the total tributary escapement estimates. If outplants successfully reproduce, future jack and adult returns to the Walla Walla Basin will need to be included in estimates of mainstem tributary escapements.

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Yakima River BRT Stock		
Total run to the river		
1954-1956	Mongillo and Falconer, 1980 as cited in Howell et al. 1985.	Interpolated from Figure 2, page 394B in Howell et al. 1985 citing Mongillo and Falconer, 1980. Assumes Y-axis origin is zero rather than the 3,000 value printed in the figure.
Jack and adult run to the river (used adults)		
1957-1959	Fast et al. Unknown date (for adult return estimates); Personal communication, B. Bosch, Yakama Nation Fisheries Resource Management, Yakima, Washington, 3/2002 (for river run estimates by adults and jacks, 1982-2001).	Adults interpolated from Figure 2, page 26 in Fast et al. Unknown date; The average jack/adult ratio for the Yakima Basin from 1982-2001 (0.123) is used to estimate jack returns.
Yakima River BRT Stock		
1960-1981	Fast et al. Unknown date; Matylewich, undated in Schaller et al. 1992; ODFW and WDFW, 2000 (for jack and adult passage counts at McNary Dam). Bosch 2001; Personal communication, B. Bosch, Yakama Nation Fisheries Resource Management, Yakima, Washington, 3/2002 (for river run estimates by adults and jacks, 1982-2001).	Adults interpolated from Figure 2, page 26 in Fast et al. Unknown date were cross referenced with numerical values in Matylewich (undated). The relative difference from both sources averages 0.005, and Matylewich (undated) is used. The jack/adult ratio for each year for McNary Dam is used to estimate Yakima Basin jacks.
1982-2001	Yakama Nation Fisheries Resource Management, Yakima, Washington, 3/2002 (for river run estimates by adults and jacks, 1982-2001).	Yakama Nation Fisheries Resource Management spring Chinook salmon run reconstruction dataset.
Snake Spring/Summer Chinook		
Tucannon River BRT Stock		
Total run to the river (used through 1978)		
1969-1984	Gallinat et al. 2001 (for redd counts and fish per redd); Schwartzberg and Roger, 1986 (for historic redd counts) and Howell et al. 1985 (for historic redd counts and harvest estimates).	For the EDT project conversion rate estimates, an approximation of escapement potential is estimated by expanding historic redd counts to other spawning areas with a factor of 6.7, and expanding total potential redds by 2.6 fish per redd. Adult harvest during open seasons (1969-1973) estimated from punch cards is added to approximate a total potential run to the river.
A consistent time series of redd counts is available for the reach from Tucannon Camp Ground to Cow Camp Bridge (except 1978 and 1979). Redd counts in this reach represent approximately 15 percent of total redds during the 2000 return year (Table 7 in Gallinat et al. 2001). The annual fish per redd ratio averages approximately 2.6 for the period 1985-2000 (Table 13 in Gallinat et al. 2001). For the two years with no redd counts, the 1977 approximate potential return was projected through 1978 and 1979 at the proportional rate of change of McNary Dam passage counts. Cautions to consider regarding temporal changes in spawning distribution are described in Gallinat et al. 2001). The estimates described here follow generally an annual 200 adult escapement level after 1971 and remain below an average 2,400 fish per year escapement prior to 1953 reported in WDF et al. (1990).		
Jack and adult run to the river		
1979-2002	Personal communication, M. Gallinat, gallimpq@dfw.wa.gov , WDFW, Dayton, 5/2003	WDFW Tucannon River spring Chinook escapement dataset.

Tributary			
Attribute			
	Time period	Information sources	Methods, assumptions, or comments
Lower Granite Dam			
Total adult count			
	1975-2001	PSC Chinook TAC data - S. Kiefer, skiefer@idfg.state.id.us , IDFG Boise, pers. comm. 05/2003	Unknown methods; generally use run timing to distinguish spring/summer/fall, length criteria to distinguish jacks and adults, and fin clips (recent) or hatchery returns (historical) plus some unknown prespawm mortality to distinguish hatchery/wild.
Wild adult count			
	1962-1978	Petrosky et al. 2001	Spawners included wild- and hatchery-origin fish spawning in tributaries. The uppermost dam changed during this era as new projects were added: Ice Harbor in 1961, Lower Monumental in 1969, Little Goose in 1970, and Lower Granite in 1975. Spawner (parent) and Dam counts used to derive spawner (S) estimates included data for brood years 1962–1997. The dam counts are total numbers of adult (ages 4–5) spring and summer Chinook for wild- and hatchery origin adults combined. The wild component was estimated by subtracting tributary harvest of wild fish and the total hatchery run size from the combined wild and hatchery dam count. Prespawning survival rate (=0.8).
	1979-2005	WDFW 2006	Generally use run timing to distinguish spring/summer/fall, length criteria to distinguish jacks and adults, and fin clips (recent) or hatchery returns (historical) plus some unknown prespawm mortality to distinguish hatchery/wild. LWG count minus estimated hatchery escapement except for 1996. To account for loss and strays, hatchery return plus harvest was divided by 0.9. In 85-86 divided by 0.8.
Salmon, Grande Ronde, and Imnaha Rivers			
Natural origin adult spawners (Index Stocks)			
	1957-1995	Beamesderfer et al. 1997 in Marmorek et al. 1998	General run reconstruction methods using, index and extensive redd counts expanded to spawners, estimates of hatchery fish on the spawning grounds.
Redd Density			
	1957-2002	Paulsen and Fisher 2002	Redd survey information (redds counted and kilometers surveyed) was obtained from various sources for the spawning streams (P. Keniry, pkeniry@eou.edu , ODFW, 107 20th Street LaGrande OR 97850, personal communication, 2002, 2003 [Grande Ronde and Imnaha basins] and Hassemer 1993; Elms-Cockrum 2001, updated by S. Keifer, skiefer@idfg.state.id.us , IDFG, 600 S Walnut St. Boise ID 83707, personal communication, 2002, 2003 [Salmon Basin]). Redds were total redds from the best available survey, from both index and non-index areas
	2003		Same methods as Paulsen and Fisher 2003
Minimum Total Spawners			
	1957-2002	Redd survey information from above sources; total redds for index and non-index areas expanded by net redd-adult ratio from Beamesderfer et al. 1997	.
BRT Stocks – Lemhi R., Pahsimeroi R., Minam R., Catherine Cr., Wenaha R.			
	Varies	Assumed from Beamesderfer et al. 1997 in Marmorek et al. 1998 for most years; post-1995 are from an unknown source.	General run reconstruction methods using, index and extensive redd counts expanded to spawners, estimates of hatchery fish on the spawning grounds.

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Snake River Fall Chinook		
Lower Granite Dam		
Total wild return over the dam		
1975-1985	BRT spreadsheet	methods not specified
1986-2003	Columbia River Compact 2004	methods not specified
Snake River Sockeye		
Various Points		
Total run		
1954-1966	BRT spreadsheet.	Redfish Lake outlet trap adult counts
1967-1974	BRT spreadsheet.	Ice Harbor Dam adult counts
1975-2002	NOAA Fisheries Biological Recovery Team. 2003	Sawtooth Hatchery trap adult counts
2003-2004	IDFG	Sawtooth Hatchery trap adult counts; 2004 is trap count + adults netted below the weir
Snake River Steelhead		
Lower Granite Dam		
A & B Run Summer Steelhead		
1980-1997	BRT spreadsheet ; CRFMP Technical Advisory Committee (TAC)	Estimates of wild adult A & B-run summer steelhead by length class
1998-2001	BRT spreadsheet ; P. Dygert, NOAA Fisheries, Montlake WA	Estimates of wild adult A & B-run summer steelhead by length class
2002-2003	S. Marshall, smarshal@idfg.state.id.us , IDFG Boise, pers. comm. 8/27/2004	Estimates of wild adult A & B-run summer steelhead by length class
Mid-Columbia River Steelhead		
BRT Stocks		
Composite adult run		
1980-2001	BRT spreadsheet	Composite of 4 BRT runs: Unknown, Yakima, Umatilla, and Touchet (from Joe Bumgarner)
Other Stocks		
Touchet River adults		
1987-2002	Joe Bumgarner, bumgajdb@dfw.wa.gov , WDFW Dayton WA, pers. comm. 05/2003	Estimates of spawning adults from trap and redd count data
Deschutes River wild adults		
1978-2002	S. Pribyl, steve.pribyl@state.or.us , ODFW, The Dalles, pers. comm. 5/2004	Table 37. Estimated number of steelhead that migrated past Sherars Falls, by run year.

PRELIMINARY ABUNDANCE-BASED TREND RESULTS FOR COLUMBIA BASIN SALMON AND STEELHEAD ESUS

Tributary		
Attribute		
Time period	Information sources	Methods, assumptions, or comments
Upper Columbia River Spring Chinook		
BRT Stock		
Composite adult run		
1960-1978	BRT spreadsheet update	Wenatchee spring Chinook (w/o Icicle Cr.) expansions from redd counts, Methow River spring Chinook, Entiat River spring Chinook
1978-2004	NMFS 2005	Methods unspecified; similar to BRT but spawners proportionally higher in years in which datasets overlap.
Upper Columbia River Summer/Fall Chinook		
Priest Rapids Dam		
Adult dam count		
1960-2003	Columbia River DART	Adult Chinook dam count for summer and fall counting periods
Upper Columbia River Steelhead		
Priest Rapids Dam		
Estimated wild adults counted at dam		
1986-2003	BRT spreadsheet update	Priest Rapids Counts; used since wild steelhead not updated by NOAA Fisheries; WDFW proportion wild since 1986 only
Lake Wenatchee Sockeye		
Columbia River turnoff		
Lake Wenatchee turnoff from the mainstem Columbia		
1960-2004	Columbia River DART	Rock Island minus Rocky Reach adults and jack count
Okanagan Sockeye		
Wells Dam		
Adults counted at dam		
1960-2004	Columbia River DART	Dam count used due to availability over BC spawning ground expansions

Table 3. NOAA Fisheries interim recovery targets (Lohn 2002), and types of data we have on hand to address progress toward those targets. Shaded targets are sums of targets for spawning aggregations within the geographic area that were in the original document.

Geographic Spawning Aggregations		Interim Abundance Targets		Data to Address Targets			
ESU/Spawning Aggregation	Index Areas	Spawning Aggregation	Index Areas	Dam Count	# Spawners	Redd Census	Redd Index
Upper Columbia Spring Chinook ESU		6,250		PRD?	Σ4 pops.?	Y?	Y
Methow River	Methow	2,000	2,000	?	BRT est.	Y	
Entiat River	Entiat	500	500		BRT est.	Y	
Okanogan River		none		Y	N	N	N
Wenatchee River	Wenatchee	3,750	3,750		BRT est.	Y	
Snake River Spring/Summer Chinook ESU		41,900		LGR	BRT/TAC est.	N	Y
Tucannon River		1,000		N	TRT est.	Y	Y
Grande Ronde River		2,000		N	N	N	Σ 9 indices
	Minam		439	N	BRT est.	Y	N
Imnaha River		2,500		N	N	N	Σ 3 indices
	Mainstem		802	N	BRT est.	N	Y
Lower Mainstem tribs.		1,000		N	N	N	Σ 2 indices
Salmon River		35,400		N	N	N	Y
Little Salmon River Basin		1,800		Rapid R. weir	N	N	N
Mainstem Salmon small tribs.		700		N	N	N	N
South Fork Salmon (Sum.)		9,200		N	Σ 2 BRT est.	N	Σ 4 indices
	Johnson Creek		288	N	BRT est.	N	Y
Middle Fork Salmon		9,300					R. Thurow, USFS IRC
	Bear Valley/Elk		911	N	BRT est.	N	Y
	Marsh Creek		426	N	BRT est.	N	Y
Mainstem Tributaries (Middle Fork to Lemhi)		700		N	N	N	N. Fk. Salmon
Lemhi River		2,200		N	BRT est.	N	Σ 2 indices
Pahsimeroi (Sum.)		1,300		Weir	Weir - Hatchery	N	Y
Mainstem tribs. (Sum.) Lemhi to Redfish Lake Creek		2,000		N	N	N	Σ 2 indices
Mainstem tribs. (Spr.) Lemhi to Yankee Fork		2,400		N	N?	N	Σ 2 indices
Upper East Fork tribs. (Spr.)		700		N	N?	N	Y
Upper Salmon Basin (Spr.)		5,100		N	Valley Cr. BRT est.	N	Σ 3-4 indices
Snake River Fall Chinook ESU		2,500		LGR	BRT est.	Y	Y
Snake River Sockeye ESU		1000 in one lake; 500 per year in second lake		Sawtooth weir	Redfish L. weir	N	N
Snake River Steelhead ESU		53,700		LGR	LGR TAC	N	Approx. 42 streams
Tucannon River		1,300		N	BRT est.	N	Y?
Asotin Creek		400		N	BRT est.	N	Y

Table 3. (Continued).

Geographic Spawning Aggregations		Interim Abundance Targets		Data to Address Targets			
ESU/Spawning Aggregation	Index Areas	Spawning Aggregation	Index Areas	Dam Count	# Spawners	Redd Census	Redd Index
Snake River Steelhead ESU (continued)							
Grande Ronde River		10,000					18 streams
Lower Grande Ronde R.		2,600		N	BRT est. (1 stream)?	N	1 stream?
Joseph Creek		1,400		N	BRT est. (9 streams)	N	9 streams
Middle Fork		2,000		N	BRT est. (3 streams?)	N	3 streams in Wallowa?
Upper Mainstem Grande Ronde R.		4,000		N	BRT est. (5 streams?)	N	5 streams?
Imnaha River		2,700		N	BRT est. (6 streams)	N	6 streams
Clearwater River		17,700		Lewiston Dam (1971)			
Mainstem Clearwater R.		4,900		N	N	N	N
South Fork Clearwater R		3,400		Crooked R Weir	N	N	N
Middle Fork Clearwater R		1,700		Fish Cr. Weir	N	N	N
Selway R.		4,900		N	N	N	N
Lochsa R.		2,800		N	N	N	N
Salmon River		21,600					
Lower Salmon R.		1,700		N	N	N	N
Little Salmon R.		1,400		N	N	N	N
South Fork Salmon R.		4,000		N	N	N	N
Middle Fork Salmon R.		7,400		N	N	N	N
Upper Salmon R.		4,700		N	N	N	N
Lemhi R.		1,600		N	N	N	N
Pahsimeroi R.		800		N	N	N	N
Upper Columbia Steelhead ESU		5,500		N	Σ BRT est.	N	2 trends
Methow River		2,500		WEL	BRT (2 trends)	N	Y
Entiat River		500		N	BRT (w/ Wenatchee)	N	N
Okanogan River		none		WEL	WDFW?	N	Y?
Wenatchee River		2,500		N	BRT (w/ Entiat)	N	Disc.
Middle Columbia Steelhead ESU		55,400		N	N	N	4 areas
Yakima River		10,500			BOR (-2000)		
Satus Cr./Toppenish Cr.		2,400		N	N	N	N
Naches R.		3,400		N	N	N	N
Mainstem Yakima R. (Wapato to Roza)		1,800		N	N	N	N
Mainstem Yakima R. (above Roza)		2,900		N	N	N	N

Table 3. (Continued).

Geographic Spawning Aggregations		Interim Abundance Targets		Data to Address Targets			
ESU/Spawning Aggregation	Index Areas	Spawning Aggregation	Index Areas	Dam Count	# Spawners	Redd Census	Redd Index
Middle Columbia Steelhead ESU (continued)					N	N	4 areas
Klickitat River		3,600		N	N	N	N
Walla-Walla River		2,600		N	N	N	N
Umatilla River		2,300		3 Mile Dam?	N	N	N
Deschutes River (below Pelton Dam complex)		6,300		Sherars Falls	N	N	Y
John Day River		9,800					
North Fork John Day R.		2,700		N	TRT est.	Y	Y
Middle Fork John Day R.		1,300		N	TRT est.	Y	Y
South Fork John Day R.		600		N	N	N	N
Lower John Day R.		3,200		N	N	N	N
Upper John Day R.		2,000		N	TRT est.	Y	Y

Table 4. Geometric mean abundance and trends for the 19 ESUs in the Columbia River basin, using a 10-year base period. Descriptions of the columns follow the table. (LCR = Lower Columbia River; SR = Snake River; MCR = Mid-Columbia River; UCR = Upper Columbia River). Note that one ESU did not have enough post-2000 data to calculate percent change or perform the BiOp test analogues.

Trend No.	ESU	Aggregation	Type	Most Recent Year	Interim Target	Geomean (1996-2000)	Geomean (2001-recent)	Trend Slope (1990-2000)	Trend Slope (1990-recent)	% change in geomean	% change in trend	Change in trend +?	Change in mean +?	Trend > 1?
1	LCR Chinook	Lower Columbia ESU	Best Total Spawners	2004		11,033	32,990	0.96	1.08	199%	11.9%	True	True	True
2	MCR Spring Chinook	Mid-Columbia ESU	Best Total Spawners	2004		13,367	35,177	1.02	1.10	163%	8.0%	True	True	True
3	SR Spring Chinook	Snake River ESU	Wild Adults	2005	41,900	4,840	25,957	0.94	1.11	436%	18.0%	True	True	True
4	SR Fall Chinook	Above Lower Granite Dam	Wild Adults	2004	2,500	695	3,748	1.16	1.25	439%	7.6%	True	True	True
5	SR Sockeye	Snake River ESU	Adults	2005	1,500	4	15	1.23	1.19	224%	-4.0%	False	True	True
6	SR Steelhead	A & B Runs above Lower Granite Dam	Wild Adults	2003	53,700	10,694	37,784	1.00	1.10	253%	9.3%	True	True	True
7	UCR Spring Chinook	Upper Columbia ESU	Adults	2004	6,250	374	2,207	0.82	1.01	490%	23.0%	True	True	True
8	UCR Steelhead	Upper Columbia River ESU	Wild Adults	2003	5,500	1,146	3,643	0.97	1.06	218%	9.2%	True	True	True
9	Deschutes R Summer/Fall Ch	Deschutes River	Wild Adults	2003		9,137	12,773	1.10	1.10	40%	0.1%	True	True	True
10	MCR Steelhead	Mid-Columbia River Steelhead ESU Composite	Wild Adults	2002	55,400	7,228	17,553	0.99	1.05	143%	6.2%	True	True	True
11	U Willamette Spring Chinook	Clackamas & McKenzie Rivers	Adults	2005		3,041	13,080	0.89	1.05	330%	18.7%	True	True	True
12	U Willamette Steelhead	Willamette Falls Dam Winter Run	Natural Adults	2006		3,961	8,288	0.93	1.02	109%	9.8%	True	True	True
13	LCR Fall Chinook	14 Early, Mid- and Late Fall Run Stocks	Adults	2000		20,698	NA	0.97	0.97	NA	0.0%	N/A	N/A	N/A
14	LCR Chum	Grays R. & Hamilton / Hardy Cr.	Adults	2003		2,114	1,776	1.02	1.00	-16%	-1.5%	False	False	True
15	LCR Steelhead	Lower Columbia ESU	Adults	2001		6,333	4,429	0.93	0.92	-30%	-0.8%	False	False	False
16	LCR Coho	Sandy & Clackamas Rivers	Adults	2005		822	2,648	0.92	1.02	222%	11.2%	True	True	True
17	L Wenatchee Sockeye	Lake Wenatchee Turnoff	Adult Escapement	2005		7,588	18,489	0.88	1.01	144%	13.7%	True	True	True
18	Okanogan R Sockeye	Okanogan R ESU	Adult Escapement	2005		17,211	39,502	1.02	1.10	130%	7.6%	True	True	True
19	UCR Summer/Fall Chinook	Upper Columbia River ESU	Adult Escapement	2005		37,530	111,144	1.09	1.14	196%	4.6%	True	True	True

¹Snake River sockeye returns of zero (1990, 1995, and 1997) were changed to 1 in order to avoid a geometric mean of zero

Table 4. (Continued).

Key for Table 4

ESU: The ESU in which the spawning aggregation resides

Aggregation: Spawning aggregation used for trend, i.e. selected stocks or rivers, entire ESU, etc.

Trend type: e.g., total spawners, wild spawners

Most recent year: Most recent year of data available for the spawning aggregation

Geomean (1996-2000): Geometric mean based on years 1996-2000

Geomean (2001-recent): Geometric mean based on years post-2000

Trend Slope (1990-2000): Trend calculation used in recent BRT document based on 1990-2000

Trend Slope (1990- recent): Trend calculation used in recent BRT document based on 1990-most recent year

N/A: Means that the indicated quantity is not available, due to lack of data

% Change in geomean: The % change in the geomean between periods 1996-2000 and 2001-recent

% Change in trend: The % change in trend between periods 1990-2000 and 1990-recent

Change in trend +?: TRUE if % Change in trend is positive, FALSE otherwise

Change in mean +?: TRUE if % Change in geomean is positive, FALSE otherwise

Trend > 1?: TRUE if BRT Trend (1990-recent) is greater than one, FALSE otherwise

Table 5. Trends for the 19 ESUs in the Columbia River basin, using a 20-year base period. Descriptions of the columns follow Table 4. (LCR = Lower Columbia River; SR = Snake River; MCR = Mid-Columbia River; UCR = Upper Columbia River). Note that one ESU did not have enough post-2000 data to calculate percent change or perform the BiOp test analogues.

Trend No.	ESU	Aggregation	Type	Most Recent Year	Trend Slope (1980-2000)	Trend Slope (1980-recent)	% change in trend	Change in trend +?	Trend > 1?
1	LCR Chinook	Lower Columbia ESU	Best Total Spawners	2004	1.04	1.07	3.0%	True	True
2	MCR Spring Chinook	Mid-Columbia ESU	Best Total Spawners	2004	1.02	1.05	3.0%	True	True
3	SR Spring Chinook	Snake River ESU	Wild Adults	2005	0.95	1.02	7.4%	True	True
4	SR Fall Chinook	Above Lower Granite Dam	Wild Adults	2004	1.03	1.09	6.0%	True	True
5	SR Sockeye	Snake River ESU	Adults	2005	0.85	0.93	9.4%	True	False
6	SR Steelhead	A & B Runs above Lower Granite Dam	Wild Adults	2003	0.96	1.00	4.0%	True	False
7	UCR Spring Chinook	Upper Columbia ESU	Adults	2004	0.87	0.93	7.2%	True	False
8	UCR Steelhead	Upper Columbia River ESU	Wild Adults	2003	0.93	0.99	7.3%	True	False
9	Deschutes R Summer/Fall Ch	Deschutes River	Wild Adults	2003	1.03	1.05	1.1%	True	True
10	MCR Steelhead	Mid-Columbia River Steelhead ESU Composite	Wild Adults	2002	0.96	0.98	2.2%	True	False
11	U Willamette Spring Chinook	Clackamas & McKenzie Rivers	Adults	2005	0.98	1.03	4.6%	True	True
12	U Willamette Steelhead	Willamette Falls Dam Winter Run	Natural Adults	2006	0.92	0.96	4.5%	True	False
13	LCR Fall Chinook	14 Early, Mid- and Late Fall Run Stocks	Adults	2000	0.98	0.98	0.0%	N/A	N/A
14	LCR Chum	Grays R. & Hamilton / Hardy Cr.	Adults	2003	1.07	1.05	-1.6%	False	True
15	LCR Steelhead	Lower Columbia ESU	Adults	2001	0.97	0.97	-0.6%	False	False
16	LCR Coho	Sandy & Clackamas Rivers	Adults	2005	0.94	0.98	4.0%	True	False
17	L Wenatchee Sockeye	Lake Wenatchee Turnoff	Adult Escapement	2005	0.92	0.96	4.8%	True	False
18	Okanogan R Sockeye	Okanogan R ESU	Adult Escapement	2005	0.94	0.99	5.3%	True	False
19	UCR Summer/Fall Chinook	Upper Columbia River ESU	Adult Escapement	2005	1.02	1.06	3.7%	True	True

¹Snake River sockeye returns of zero (1990, 1995, and 1997) were changed to 1 in order to avoid a geometric mean of zero

Table 6. Geometric mean abundance and trends for 68 of the individual populations in the Columbia River basin, using a 10-year base period. Descriptions of the columns follow Table 4. (LCR = Lower Columbia River; SR = Snake River; MCR = Mid-Columbia River; UCR = Upper Columbia River).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Interim Target	Geomean (1996-2000)	Geomean (2001-recent)	Trend Slope (1990-2000)	Trend Slope (1990-recent)	% change in geomean	% change in trend	Change in trend +?	Change in mean +?	Trend > 1?
1	LCR Chinook	Hood River Spring Chinook	Best Total Spawners	2004		153	513	0.76	0.92	236%	21%	True	True	False
2	LCR Chinook	Little White Salmon River Spring Chinook	Adults	2004		3,985	12,976	1.03	1.13	226%	10%	True	True	True
3	LCR Chinook	White Salmon River Spring Chinook	Adults	2004		177	339	0.78	0.90	92%	15%	True	True	False
4	LCR Chinook	Wind River Spring Chinook	Best Total Spawners	2004		6,459	18,624	0.96	1.07	188%	12%	True	True	True
5	MCR Steelhead	Deschutes River	Wild Adults	2003	6,300	2,857	9,029	1.01	1.11	216%	10%	True	True	True
6	MCR Steelhead	Deschutes River Eastside	Total Adults	2005		1,896	4,650	1.06	1.11	145%	4%	True	True	True
7	MCR Steelhead	Deschutes River Westside	Total Adults	2005		433	948	1.05	1.09	119%	4%	True	True	True
8	MCR Steelhead	Fifteenmile Creek	Total Adults	2005		419	840	1.07	1.11	101%	4%	True	True	True
9	MCR Steelhead	John Day River Lower Mainstem	Spawners	2005	3,200	1,523	2,628	1.04	1.05	73%	1%	True	True	True
10	MCR Steelhead	John Day River Middle Fork	Spawners	2005	1,300	684	995	0.96	0.99	45%	3%	True	True	False
11	MCR Steelhead	John Day River North Fork	Spawners	2005	2,700	1,536	2,345	1.13	1.10	53%	-2%	False	True	True
12	MCR Steelhead	John Day River South Fork	Spawners	2005	600	156	515	0.89	1.02	231%	14%	True	True	True
13	MCR Steelhead	John Day River Upper Mainstem	Spawners	2005	2,000	462	707	0.90	0.97	53%	8%	True	True	False
14	MCR Steelhead	Touchet River	Total Wild Fish	2004		303	201	0.99	0.96	-34%	-3%	False	False	False
15	MCR Steelhead	Umatilla River	Total Adults	2004	2,300	1,913	3,512 ¹	1.07	1.09	84%	3%	True	True	True
16	MCR Steelhead	Walla Walla River	Total Adults	2003	2,600	750	1,741	0.92	1.04	132%	13%	True	True	True
17	MCR Steelhead	Yakima River Naches River	Total Adults	2004	3,400	334	1,001	0.97	1.07	199%	10%	True	True	True
18	MCR Steelhead	Yakima River Satus Creek	Total Adults	2004		411	1,230	0.98	1.07	199%	9%	True	True	True
19	MCR Steelhead	Yakima River Toppenish Creek	Total Adults	2004	2,400	107	321	0.99	1.08	199%	9%	True	True	True

¹ It is unknown whether the spawning aggregation meets the NOAA Fisheries interim target since the count includes hatchery adults.

Table 6. (Continued).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Interim Target	Geomean (1996-2000)	Geomean (2001-recent)	Trend Slope (1990-2000)	Trend Slope (1990-recent)	% change in geomean	% change in trend	Change in trend +?	Change in mean +?	Trend > 1?
20	MCR Steelhead	Yakima River Upper Yakima River	Total Adults	2004	2,900	58	200	0.98	1.09	244%	12%	True	True	True
21	MCR Spring Chinook	Deschutes River Spring Chinook	Adults	2004		3,383	8,896	0.97	1.07	163%	10%	True	True	True
22	MCR Spring Chinook	Deschutes River Spring Chinook	Wild Adults	2003		857	1,799	0.92	1.01	110%	10%	True	True	True
23	MCR Spring Chinook	John Day River Middle Fork Spring Chinook	Total Adult Spawners	2003		560	963	1.13	1.14	72%	1%	True	True	True
24	MCR Spring Chinook	John Day River North Fork Spring Chinook	Total Adult Spawners	2003		1,266	2,900	0.99	1.06	129%	7%	True	True	True
25	MCR Spring Chinook	John Day River Spring Chinook	Best Total	2004		2,592	5,201	1.03	1.09	101%	5%	True	True	True
26	MCR Spring Chinook	John Day River Upper Spring Chinook	Total Adult Spawners	2003		439	1,129	1.03	1.10	157%	7%	True	True	True
27	MCR Spring Chinook	Klickitat River Spring Chinook	Best Total	2004		1,244	2,447	0.95	1.03	97%	9%	True	True	True
28	MCR Spring Chinook	Umatilla River Spring Chinook	Adults	2004		1,738	4,413	1.08	1.15	154%	7%	True	True	True
29	MCR Spring Chinook	Yakima River Spring Chinook	Best Total	2006		3,971	9,890	1.08	1.12	149%	3%	True	True	True
30	SR Spring Chinook	Bear Valley Creek Chinook	Natural Run	2003	911	162	1,001	1.00	1.16	518%	16%	True	True	True
31	SR Spring Chinook	Big Creek Chinook	Natural Run	2004		45	414	0.94	1.14	818%	21%	True	True	True
32	SR Spring Chinook	Camas Creek Chinook	Natural Run	2004		13	175	0.97	1.20	1282%	24%	True	True	True
33	SR Spring Chinook	Catherine Creek Chinook	Total Natural Spawners	2005		61	115	1.31	1.19	89%	-9%	False	True	True
34	SR Spring Chinook	Grande Ronde Upper Mainstem Chinook	Natural Spawners	2005		35	48	0.94	1.00	37%	6%	True	True	False
35	SR Spring Chinook	Imnaha River Chinook	Total Natural Spawners	2005	802	253	836	1.05	1.11	230%	6%	True	True	True
36	SR Spring Chinook	Lemhi River Chinook	Natural Run	2003	2,200	88	249	1.04	1.12	183%	8%	True	True	True
37	SR Spring Chinook	Loon Creek Chinook	Natural Run	2004		22	354	1.13	1.31	1502%	15%	True	True	True
38	SR Spring Chinook	Lostine River Chinook	Natural Adults	2005		168	423	1.13	1.16	152%	2%	True	True	True

¹ It is unknown whether the spawning aggregation meets the NOAA Fisheries interim target since the count includes hatchery adults.

Table 6. (Continued).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Interim Target	Geomean (1996-2000)	Geomean (2001-recent)	Trend Slope (1990-2000)	Trend Slope (1990-recent)	% change in geomean	% change in trend	Change in trend +?	Change in mean +?	Trend > 1?
39	SR Spring Chinook	Lower Mainstem Salmon River	Natural Run	2005		69	221	0.99	1.11	217%	11%	True	True	True
40	SR Spring Chinook	Marsh Creek Chinook	Natural Run	2003	426	25	412	0.85	1.11	1540%	31%	True	True	True
41	SR Spring Chinook	Minam River Chinook	Total Natural Spawners	2005		247	421	1.12	1.13	70%	1%	True	True	True
42	SR Spring Chinook	Secesh River Chinook	Natural Run	2003		252	1,024	1.01	1.12	305%	11%	True	True	True
43	SR Spring Chinook	Snake River Spring/Summer Chinook ESU	Total Adults	2005		16,063	83,642	1.00	1.15	421%	15%	True	True	True
44	SR Spring Chinook	South Fork Salmon East Fork (incl. Johnson Cr.)	Natural Run	2003	288	257	1,045 ¹	0.88	1.03	307%	17%	True	True	True
45	SR Spring Chinook	South Fork Salmon Mainstem	Natural Run	2001		789	1,446	0.98	1.01	83%	3%	True	True	True
46	SR Spring Chinook	Sulphur Creek	Natural Run	2003		18	142	0.78	1.00	703%	28%	True	True	True
47	SR Spring Chinook	Tucannon River Spring Chinook	Adults	2005	1,000	56	588	0.75	1.04	954%	38%	True	True	True
48	SR Spring Chinook	Tucannon Spring Chinook	Natural Run	2003	1,000	144	445	0.87	0.99	209%	14%	True	True	False
49	SR Spring Chinook	Upper Mainstem Salmon River	Natural Run	2005		151	477	0.90	1.05	215%	16%	True	True	True
50	SR Spring Chinook	Valley Creek Chinook	Natural Run	2003		23	246	1.00	1.20	955%	21%	True	True	True
51	SR Spring Chinook	Wenaha River Chinook	Total Natural Spawners	2005		247	601	1.17	1.20	144%	2%	True	True	True
52	SR Spring Chinook	Yankee Fork Salmon River	Natural Run	2003		8	112	0.88	1.12	1275%	27%	True	True	True
53	SR Steelhead	Asotin Creek A-Run	Total Wild Fish	2001	400	211	543	0.99	1.04	158%	5%	True	True	True
54	SR Steelhead	Grande Ronde River Upper Mainstem A-Run	Expanded Redds	2004	4,000	2,435	1,877	1.03	1.00	-23%	-3%	False	False	False
55	SR Steelhead	Grande Ronde Joseph Cr A-Run	Expanded Redds	2005	1,400	1,260	1,917	1.01	1.03	52%	3%	True	True	True
56	SR Steelhead	Grande Ronde Whisky & Prairie Creeks A-Run	Expanded Redds	2005		114	141	1.08	1.07	24%	-2%	False	True	True
57	SR Steelhead	Imnaha Camp Cr A-Run	Expanded Redds	2005		45	105	0.96	1.05	134%	9%	True	True	True

¹ It is unknown whether the spawning aggregation meets the NOAA Fisheries interim target since the count includes hatchery adults.

Table 6. (Continued).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Interim Target	Geomean (1996-2000)	Geomean (2001-recent)	Trend Slope (1990-2000)	Trend Slope (1990-recent)	% change in geomean	% change in trend	Change in trend +?	Change in mean +?	Trend > 1?
58	SR Steelhead	Imnaha Little Sheep Cr A-Run	Total Wild Fish	2002		20	136	0.97	1.09	564%	12%	True	True	True
59	SR Steelhead	Snake River Summer Steelhead A-Run	Total Wild Fish	2003	53,700	8,884	29,421	1.03	1.11	231%	8%	True	True	True
60	SR Steelhead	Snake River Summer Steelhead B-Run	Total Wild Fish	2003		1,672	6,915	0.91	1.04	314%	15%	True	True	True
61	SR Steelhead	Tucannon River A-Run	Total Wild Fish	2001	1,300	67	198	0.85	0.91	195%	7%	True	True	False
62	UCR Steelhead	Entiat River	Total Spawners	2004	500	280	593 ¹	0.98	1.05	112%	7%	True	True	True
63	UCR Steelhead	Methow River	Total Spawners	2005	2,500	1,280	5,936 ¹	1.01	1.14	364%	12%	True	True	True
64	UCR Steelhead	Okanogan River	Total Spawners	2005		660	3,106	1.01	1.14	370%	12%	True	True	True
65	UCR Steelhead	Wenatchee River	Total Spawners	2004	2,500	2,667 ¹	5,648 ¹	0.98	1.05	112%	7%	True	True	True
66	UCR Spring Chinook	Entiat River	Total Wild Fish	2004	500	65	263	0.94	1.08	308%	15%	True	True	True
67	UCR Spring Chinook	Methow River	Total Wild Fish	2004	2,000	26	575	0.68	0.94	2143%	37%	True	True	False
68	UCR Spring Chinook	Wenatchee River	Total Wild Fish	2004	3,750	214	1,148	0.84	1.03	437%	21%	True	True	True

¹ It is unknown whether the spawning aggregation meets the NOAA Fisheries interim target since the count includes hatchery adults.

Table 7. Geometric mean abundance and trends for 68 of the individual populations in the Columbia River basin, using a 20-year base period. Descriptions of the columns follow Table 4. (LCR = Lower Columbia River; SR = Snake River; MCR = Mid-Columbia River; UCR = Upper Columbia River).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Trend Slope (1980-2000)	Trend Slope (1980-recent)	% change in trend	Change in trend +?	Trend > 1?
1	LCR Chinook	Hood River Spring Chinook	Best Total Spawners	2004	0.85	0.96	14%	True	False
2	LCR Chinook	Little White Salmon River Spring Chinook	Adults	2004	1.04	1.08	4%	True	True
3	LCR Chinook	White Salmon River Spring Chinook	Adults	2004	1.41	1.31	-7%	False	True
4	LCR Chinook	Wind River Spring Chinook	Best Total Spawners	2004	1.04	1.07	3%	True	True
5	MCR Steelhead	Deschutes River	Wild Adults	2003	0.94	0.98	5%	True	False
6	MCR Steelhead	Deschutes River Eastside	Total Adults	2005	1.06	1.11	4%	True	True
7	MCR Steelhead	Deschutes River Westside	Total Adults	2005	0.97	1.00	4%	True	True
8	MCR Steelhead	Fifteenmile Creek	Total Adults	2005	0.99	1.04	6%	True	True
9	MCR Steelhead	John Day River Lower Mainstem	Spawners	2005	0.96	0.98	3%	True	False
10	MCR Steelhead	John Day River Middle Fork	Spawners	2005	0.96	0.97	1%	True	False
11	MCR Steelhead	John Day River North Fork	Spawners	2005	0.96	1.00	3%	True	False
12	MCR Steelhead	John Day River South Fork	Spawners	2005	0.91	0.95	5%	True	False
13	MCR Steelhead	John Day River Upper Mainstem	Spawners	2005	0.94	0.96	2%	True	False
14	MCR Steelhead	Touchet River	Total Wild Fish	2004	0.98	0.96	-2%	False	False
15	MCR Steelhead	Umatilla River	Total Adults	2004	1.01	1.03	2%	True	True
16	MCR Steelhead	Walla Walla River	Total Adults	2003	0.92	1.04	13%	True	True
17	MCR Steelhead	Yakima River Naches River	Total Adults	2004	0.93	1.00	7%	True	True
18	MCR Steelhead	Yakima River Satus Creek	Total Adults	2004	0.94	1.00	7%	True	True
19	MCR Steelhead	Yakima River Toppenish Creek	Total Adults	2004	0.94	1.01	7%	True	True
20	MCR Steelhead	Yakima River Upper Yakima River	Total Adults	2004	0.92	1.00	9%	True	False
21	MCR Spring Chinook	Deschutes River Spring Chinook	Adults	2004	1.00	1.03	3%	True	True
22	MCR Spring Chinook	Deschutes River Spring Chinook	Wild Adults	2003	0.94	0.97	3%	True	False
23	MCR Spring Chinook	John Day River Middle Fork Spring Chinook	Total Adult Spawners	2003	1.02	1.04	2%	True	True
24	MCR Spring Chinook	John Day River North Fork Spring Chinook	Total Adult Spawners	2003	1.02	1.04	2%	True	True
25	MCR Spring Chinook	John Day River Spring Chinook	Best Total	2004	1.03	1.05	2%	True	True
26	MCR Spring Chinook	John Day River Upper Spring Chinook	Total Adult Spawners	2003	1.04	1.07	2%	True	True
27	MCR Spring Chinook	Klickitat River Spring Chinook	Best Total	2004	0.97	1.00	3%	True	False
28	MCR Spring Chinook	Umatilla River Spring Chinook	Adults	2004	1.64	1.55	-6%	False	True
29	MCR Spring Chinook	Yakima River Spring Chinook	Best Total	2006	1.03	1.06	3%	True	True
30	SR Spring Chinook	Bear Valley Creek Chinook	Natural Run	2003	1.00	1.05	5%	True	True
31	SR Spring Chinook	Big Creek Chinook	Natural Run	2004	0.95	1.02	7%	True	True
32	SR Spring Chinook	Camas Creek Chinook	Natural Run	2004	0.89	0.99	11%	True	False

Table 7. (Continued).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Trend Slope (1980-2000)	Trend Slope (1980-recent)	% change in trend	Change in trend +?	Trend > 1?
33	SR Spring Chinook	Catherine Creek Chinook	Total Natural Spawners	2005	0.86	0.92	8%	True	False
34	SR Spring Chinook	Grande Ronde Upper Mainstem Chinook	Natural Spawners	2005	0.89	0.93	5%	True	False
35	SR Spring Chinook	Imnaha River Chinook	Total Natural Spawners	2005	0.93	0.98	6%	True	False
36	SR Spring Chinook	Lemhi River Chinook	Natural Run	2003	0.94	0.98	5%	True	False
37	SR Spring Chinook	Loon Creek Chinook	Natural Run	2004	0.97	1.07	10%	True	True
38	SR Spring Chinook	Lostine River Chinook	Natural Adults	2005	0.95	1.01	6%	True	True
39	SR Spring Chinook	Lower Mainstem Salmon River	Natural Run	2005	0.95	1.00	6%	True	True
40	SR Spring Chinook	Marsh Creek Chinook	Natural Run	2003	0.92	1.00	9%	True	False
41	SR Spring Chinook	Minam River Chinook	Total Natural Spawners	2005	1.03	1.06	3%	True	True
42	SR Spring Chinook	Secesh River Chinook	Natural Run	2003	0.98	1.02	4%	True	True
43	SR Spring Chinook	Snake River Spring/Summer Chinook ESU	Total Adults	2005	0.98	1.05	7%	True	True
44	SR Spring Chinook	South Fork Salmon East Fork (incl. Johnson Cr.)	Natural Run	2003	0.99	1.03	4%	True	True
45	SR Spring Chinook	South Fork Salmon Mainstem	Natural Run	2001	1.04	1.05	0%	True	True
46	SR Spring Chinook	Sulphur Creek	Natural Run	2003	0.96	1.01	5%	True	True
47	SR Spring Chinook	Tucannon River Spring Chinook	Adults	2005	0.85	0.95	11%	True	False
48	SR Spring Chinook	Tucannon Spring Chinook	Natural Run	2003	0.91	0.95	4%	True	False
49	SR Spring Chinook	Upper Mainstem Salmon River	Natural Run	2005	0.92	0.97	6%	True	False
50	SR Spring Chinook	Valley Creek Chinook	Natural Run	2003	0.94	1.02	8%	True	True
51	SR Spring Chinook	Wenaha River Chinook	Total Natural Spawners	2005	0.98	1.04	6%	True	True
52	SR Spring Chinook	Yankee Fork Salmon River	Natural Run	2003	0.95	1.03	8%	True	True
53	SR Steelhead	Asotin Creek A-Run	Total Wild Fish	2001	0.92	0.95	3%	True	False
54	SR Steelhead	Grande Ronde River Upper Mainstem A-Run	Expanded Redds	2004	1.00	0.99	-1%	False	False
55	SR Steelhead	Grande Ronde Joseph Cr A-Run	Expanded Redds	2005	0.99	1.00	1%	True	False
56	SR Steelhead	Grande Ronde Whisky & Prairie Creeks A-Run	Expanded Redds	2005	1.03	1.03	1%	True	True
57	SR Steelhead	Imnaha Camp Cr A-Run	Expanded Redds	2005	1.00	1.03	2%	True	True
58	SR Steelhead	Imnaha Little Sheep Cr A-Run	Total Wild Fish	2002	0.95	1.02	7%	True	True
59	SR Steelhead	Snake River Summer Steelhead A-Run	Total Wild Fish	2003	0.95	1.01	7%	True	True
60	SR Steelhead	Snake River Summer Steelhead B-Run	Total Wild Fish	2003	0.89	0.97	9%	True	False
61	SR Steelhead	Tucannon River A-Run	Total Wild Fish	2001	0.84	0.88	5%	True	False
62	UCR Steelhead	Entiat River	Total Spawners	2004	0.98	1.00	2%	True	True
63	UCR Steelhead	Methow River	Total Spawners	2005	1.00	1.05	5%	True	True
64	UCR Steelhead	Okanogan River	Total Spawners	2005	0.99	1.05	5%	True	True
65	UCR Steelhead	Wenatchee River	Total Spawners	2004	0.98	1.00	2%	True	True

Table 7. (Continued).

Trend No.	ESU	Aggregation	Type	Most Recent Year	Trend Slope (1980-2000)	Trend Slope (1980-recent)	% change in trend	Change in trend +?	Trend > 1?
66	UCR Spring Chinook	Entiat River	Total Wild Fish	2004	0.90	0.97	7%	True	False
67	UCR Spring Chinook	Methow River	Total Wild Fish	2004	0.85	0.92	8%	True	False
68	UCR Spring Chinook	Wenatchee River	Total Wild Fish	2004	0.85	0.92	8%	True	False

6 Figures

Figure 1. Columbia River Basin Chinook salmon ESUs. Numbers correspond to the trend numbers in [Table 4](#).




Figure 2. Columbia River Basin steelhead ESUs. Numbers correspond to the trend numbers in [Table 4](#).




Figure 3. Columbia River Basin coho salmon ESU. Numbers correspond to the trend numbers in [Table 4](#).




Figure 4. Columbia River Basin sockeye salmon ESUs. Numbers correspond to the trend numbers in [Table 4](#).




Figure 5. Columbia River Basin chum salmon ESU. Numbers correspond to the trend numbers in [Table 4](#).




Figure 6. The relationship between redd density of Snake River spring/summer Chinook (all redd surveys above Lower Granite Dam) and estimated wild adult spring/summer Chinook counted at the dam from 1962 through 2004. The regression is in [Appendix E](#).




Figure 7. Abundance and 5 year geometric mean of abundance for the trends best representing each of the 19 ESUs. NOAA Fisheries interim recovery targets (if any were set) are also shown. Trends prior to the 1990 BiOp baseline are shown for informational purposes.




Figure 7. (Continued).









Figure 7. (Continued).




Figure 7. (Continued).




Figure 8. Percent change in mean abundance for 18 ESU trends from the base period (5 year geometric mean of 1996 – 2000) to the test period (2001 through the most recent year). We did not have sufficient post-2000 data to calculate the change for the LCR Fall Chinook ESU.




Figure 9. Percent change in the BRT trend for 18 ESU trends from the 10-year base period (1990 – 2000) to the test period (1990 through the most recent year). We did not have post-2000 data to calculate the change for the LCR Fall Chinook ESU.





Figure 10. Percent change in the BRT trend for 18 ESU trends from the 20-year base period (1980 – 2000) to the test period (1980 through the most recent year). We did not have post-2000 data to calculate the change for the LCR Fall Chinook ESU.



7 References

(All appendices and reference documents not available on the WWW are available in a large ZIP file. Please email Tim@FisherFisheries.com for a copy of the appendices and reference documents.)

- Anonymous. 1998/1999 final hatchery escapement and broodstock report. 2000. Washington Department of Fish and Wildlife.
- Beamesderfer, R. C., H. A. Schaller, M. P. Zimmerman, C. E. Petrosky, O. P. Langness, and L. LaVoy. 1997. [Spawner-recruit data for spring and summer Chinook salmon populations in Idaho, Oregon, and Washington](#). In D. R. Marmorek and C. Peters (eds.) [Plan for analyzing and testing hypotheses \(PATH\): report of retrospective analysis for fiscal year 1997](#). ESSA Technologies Ltd, Vancouver, B.C., Canada.
- Berg, L. (editor). 2001. [Draft Yakima subbasin summary](#) prepared for the Northwest Power Planning Council through the facilities of the Columbia Basin Fish and Wildlife Authority. Portland, Oregon.
- Bosch, B. 2001. Run size forecast for Yakima River adult spring Chinook, 2002. Yakima Klickitat Fisheries Project, Yakama Nation Fisheries Resource Management. Yakima, Washington.
- Coccoli, H. (lead writer). 2001. [Draft Hood River subbasin summary](#) (including Oregon tributaries between Bonneville Dam and the Hood River) prepared for the Northwest Power Planning Council through the facilities of the Columbia Basin Fish and Wildlife Authority. Portland, Oregon.
- Columbia River Compact. 2004. [Joint Staff Report Concerning The 2004 Fall In-River Commercial Harvest Of Columbia River Fall Chinook Salmon, Summer Steelhead, Coho Salmon, Chum Salmon, And Sturgeon](#). Joint Columbia River Management Staff, ODFW and WDFW, July 21, 2004.
- CTUIR and ODFW (Confederated Tribes of the Umatilla Indian Reservation and Oregon Department of Fish and Wildlife). 1990a. [Umatilla River subbasin Salmon and Steelhead Production Plan](#), Columbia Basin System Planning. Northwest Power Planning Council and Columbia Basin Fish and Wildlife Authority.
- CTUIR and ODFW (Confederated Tribes of the Umatilla Indian Reservation and Oregon Department of Fish and Wildlife). 1990b. Walla [Walla River subbasin Salmon and Steelhead Production Plan](#), Columbia Basin System Planning. Northwest Power Planning Council and Columbia Basin Fish and Wildlife Authority.
- Daigneault, M. J., R. C. P. Beamesderfer, and G. Norman. 2003. Run Reconstructions of Select Salmon and Steelhead Populations in Washington Tributaries of the Lower Columbia River. Volume VI, Ch. 2 in [Lower Columbia Salmon and Steelhead Recovery and Subbasin Plan](#).
- Fast, D.E., Hubble, J.D. and B.D. Watson. Unknown Date. Yakima River spring Chinook - The decline and recovery of a Mid-Columbia natural spawning stock, page 18 in unknown document. Fisheries Resource Management, Yakima Indian Nation. Toppenish, Washington.
- Gallinat, M.P., J. Bumgarner, L. Ross, and M. Varney. 2001. [Tucannon River spring Chinook salmon hatchery evaluation program 2000 annual report](#). Washington Department of Fish and Wildlife, Fish Program, Science Division, Olympia, WA.
- Gephart, L. and D. Nordheim (editors). 2001. [Draft Tucannon River subbasin summary](#) prepared for the Northwest Power Planning Council through the facilities of the Columbia Basin Fish and Wildlife Authority. Portland, Oregon.

- Hammer, S., Markey, S., Adicks, K., Mains, C., Henry, M., and H. Beltran. 2001. 1999/2000 Final hatchery escapement and broodstock report. Washington Department and Fish and Wildlife, Hatcheries Division. Olympia, Washington.
- Howell, P., K. Jones, D. Scarnecchia, L LaVoy, W. Kendra, and D. Ortman. 1985. Stock assessment of Columbia River anadromous salmonids, volume 1. Report of the Oregon Department of Fish and Wildlife to Bonneville Power Administration, Portland.
- Jonasson, B. and V. Albaladejo. 1999. 1998 John Day Basin spring Chinook spawning ground survey results. Oregon Department of Fish and Wildlife memorandum, La Grande, Oregon.
- Letter from B. Lohn, NOAA Fisheries, to F. Cassidy, NPPC, April 4, 2002. Re: [Interim abundance and productivity targets for interior Columbia basin salmon and steelhead listed under the Endangered Species Act](#). Montlake, WA.
- Lindsay, R. B., B. C. Jonasson, R. K. Schroeder, and B. C. Cates. 1989. Spring Chinook salmon in the Deschutes River, Oregon. Report of Oregon Department of Fish and Wildlife to Bonneville Power Administration, Portland.
- Lindsay, R. B., W. J. Knox, M. W. Flesher, B. J. Smith, E. A. Olsen, and W. S. Lutz. 1986. Study of wild spring Chinook in the John Day River System. Report of Oregon Department of Fish and Wildlife to Bonneville Power Administration, Portland.
- Marmorek, D.R. and C.N. Peters (eds.). J. Anderson, R. Beamesderfer, L. Botsford, J. Collie, B. Dennis, R. Deriso, C. Ebbesmeyer, T. Fisher, R. Hinrichsen, M. Jones, O. Langness, L. LaVoy, G. Matthews, C. Paulsen, C. Petrosky, S. Saila, H. Schaller, C. Toole, C. Walters, E. Weber, P. Wilson, M.P. Zimmerman. 1998. [Plan for Analyzing and Testing Hypotheses \(PATH\): Retrospective and Prospective Analyses of Spring/Summer Chinook Reviewed in FY 1997](#). Compiled and edited by ESSA Technologies Ltd., Vancouver, B.C.
- Matylewich, M. Undated. Appendix B, adult dam conversion data in Schaller et al. 1992. Columbia River Intertribal Fish Commission. Portland, Oregon.
- Mongillo, P. and L. Falconer. 1980. Yakima fisheries enhancement study. Final Report. Washington Department of Game.
- Newton, J. A. 1973. Deschutes River spring Chinook salmon; a literature review. Oregon Department of Fish and Wildlife Central Region Administrative Report 74-2.
- NOAA Fisheries Biological Recovery Team. 2003. [Preliminary conclusions regarding the updated status of listed ESUs of West Coast salmon and steelhead](#). URL: <http://www.nwr.noaa.gov/BRTdraftreport/BRTdraftreport.html>
- NMFS Sustainable Fisheries Division. 2005. [Biological Opinion on Impacts of Treaty Indian and Non-Indian Fisheries in the Columbia River Basin in Years 2005-2007, on Salmon and Steelhead Listed Under the Endangered Species Act, Conference on Lower Columbia Coho, and Magnuson-Stevens Act Essential Fish Habitat Consultation](#). Seattle WA. URL: http://seahorse.nmfs.noaa.gov/pls/pcts-pub/sxn7.pcts_upload.summary_list_biop?p_id=23037
- ODFW and CTWSRO. 1990. [Hood River Subbasin Salmon and Steelhead Production Plan](#). Oregon Department of Fish and Wildlife and the Confederated Tribes of the Warm Springs Reservation. September 1990. Columbia Basin system planning. Northwest Power Planning Council and Columbia Basin Fish and Wildlife Authority.
- ODFW and WDFW. 2000. [Status report Columbia River fish runs and fisheries 1938-1999](#). Oregon Department of Fish and Wildlife, Portland, Oregon. Washington Department of Fish and Wildlife, Olympia, Washington.
- ODFW. 2001. Mid Columbia District annual Report. Oregon Department of Fish and Wildlife, The Dalles, Oregon.

- ODFW. 2004. Fisheries Management and Evaluation for 2003 Willamette River Spring Chinook. Portland, OR. URL: http://www.nwr.noaa.gov/1fmep/inplace/AnnRpts/AnnRpt2003_WillSpChFMEP.pdf
- Olsen, E .S. and R.A. French. 1996. Hood River and Pelton ladder evaluation studies. Annual Progress Report to BPA, Portland, Oregon. Oregon Department of Fish and Wildlife, Portland Oregon.
- Olsen, E. S. and R.A. French. 1998. [Hood River and Pelton ladder evaluation studies](#). Annual Progress Report to BPA, Portland, Oregon. Oregon Department of Fish and Wildlife, Portland Oregon.
- Olsen, E. S., R.A. French and A. D. Ritchy. 1994. Hood River and Pelton ladder evaluation studies. Annual Progress Report to BPA, Portland, Oregon. Oregon Department of Fish and Wildlife, Portland Oregon.
- Paulsen, C.M., and T. R. Fisher. 2003. Detecting juvenile survival effects of habitat actions: power analysis applied to endangered Snake River spring–summer Chinook (*Oncorhynchus tshawytscha*). *Can. J. Fish. Aquat. Sci.* 60: 1122-1132.
- Petit, R. 2002. Escapement database for spring Chinook in Washington tributaries between Bonneville and McNary dams, 1970-2001. Washington Department of Fish and Wildlife, Region 5. Vancouver, Washington.
- Petrosky, C. E., H. A. Schaller, and P. Budy. 2001. [Productivity and survival rate trends in the freshwater spawning and rearing stage of Snake River Chinook salmon \(*Oncorhynchus tshawytscha*\)](#). *Can. J. Fish. Aquat. Sci.* 58: 1196–1207.
- Portland General Electric. 2004. [Clackamas River Daily Fish Counts](#). Portland, OR.
- Pribyl, S. 2001. Memorandum to M. Gauvin, D. Olson, D. Ratliff, M. Paiya, September 21, 2001 - Sherars Falls harvest data, April 16, 2001 to June 15, 2001. ODFW The Dalles District Office, The Dalles, Oregon.
- Saul, D. C. Rabe, A. Davidson, and D. Rollins (lead writers). 2001. [Draft Walla Walla River subbasin summary](#) prepared for the Northwest Power Planning Council through the facilities of the Columbia Basin Fish and Wildlife Authority. Portland, Oregon.
- Schaller, H., Petrosky, C., Weber, E., and T. Cooney. 1992. Draft documentation: Chinook analytical framework Snake River spring/summer Chinook empirical life-cycle simulation model for recovery and rebuilding plan evaluation. Oregon Department of Fish and Wildlife, Portland, Oregon. Idaho Department of Fish and Game, Boise, Idaho. Columbia River Intertribal Fish Commission, Portland, Oregon. Washington Department of Fisheries, Olympia, Washington.
- Schwartzberg, M., and P. B. Roger. 1986. An annotated compendium of spawning ground surveys in the Columbia River basin above Bonneville Dam, 1960-1984. Columbia River Inter-Tribal Fish Commission, Technical Report 86-1, Portland.
- Tony, R., and R. Foster. 1978. Hatchery returns and spawning data for Columbia River, 1960-1976, progress report no. 61. Washington Department of Fisheries, Salmon Culture Division. Location unknown, Washington.
- U.S. Army Corps of Engineers. 2000. Annual fish passage - 1999. Columbia River projects - Snake River projects United States Army Corps of Engineers, North Pacific Division. Portland, Oregon.
- U.S. versus Oregon, Technical Advisory Committee. 1997. 1996 all species review, Columbia River fish management plan. Columbia River Inter-Tribal Fisheries Commission. Portland, Oregon.
- U.S. versus Oregon, Technical Advisory Committee. 2002. Spring Chinook biological assessment tables.

U.S. versus Oregon, Technical Advisory Committee. 2002. Summer Chinook biological assessment tables.

WDF, CTUIR, NPT, and WDW. 1990. [Tucannon River Subbasin Salmon and Steelhead Production Plan](#). Washington Department of Fisheries, Olympia, Washington. Confederated Tribes of the Umatilla Indian Reservation, Pendleton, Oregon. Nez Perce Tribe of Idaho, Lapwai, Idaho. Washington Department of Wildlife, Olympia, WA. September 1990. Columbia Basin system planning. Northwest Power Planning Council and Columbia Basin Fish and Wildlife Authority.

WDFW. 2004. [Lower Columbia Chum \(Grays/Sea Resources\) HGMP](#).

Wilson, W.H., J.R. Ruzycki, R.W. Carmichael, S. Onjukka, G. Claire, and J. Seals. 2001. [John Day Basin spring Chinook salmon escapement and productivity monitoring](#). Annual report to Bonneville Power Administration, Contract No. 98111646, Portland, Oregon.

8 Appendices

(All appendices and reference documents not available on the WWW are available in a 16 MB ZIP file. Please email Tim@FisherFisheries.com for a copy of the appendices and reference documents.)

- APPENDIX A. [Interim abundance and productivity targets for interior Columbia basin salmon and steelhead listed under the Endangered Species Act.](#)
- APPENDIX B. [NOAA Fisheries BRT run reconstruction data](#)
- APPENDIX C. [NOAA Fisheries Willamette/Lower Columbia Technical Recovery Team Trend Data](#)
- APPENDIX D. [Run Reconstructions For The Ecosystem Diagnosis And Treatment Model](#)
- APPENDIX E. [Various Trend Data](#)
- APPENDIX F. [Escapement trends used in the tests](#)
- APPENDIX G. [Charts for escapement trends, 5 year geometric means, and interim targets](#)